



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The dynamics of the primordial follicle reserve

Citation for published version:

Kerr, JB, Myers, M & Anderson, RA 2013, 'The dynamics of the primordial follicle reserve', *Reproduction*, vol. 146, no. 6, pp. R205-15. <https://doi.org/10.1530/REP-13-0181>

Digital Object Identifier (DOI):

[10.1530/REP-13-0181](https://doi.org/10.1530/REP-13-0181)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Reproduction

Publisher Rights Statement:

Disclaimer. This is not the definitive version of record of this article. This manuscript has been accepted for publication in [insert name of journal], but the version presented here has not yet been copy edited, formatted or proofed. Consequently, Bioscientifica accepts no responsibility for any errors or omissions it may contain. The definitive version is now freely available at <http://www.reproduction-online.org/content/146/6/R205>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The dynamics of the primordial follicle reserve

Jeffrey B Kerr^{1,2}, Michelle Myers^{1,2} and Richard A Anderson³

¹Department of Anatomy & Developmental Biology, School of Biomedical Sciences, Monash University, Victoria 3800 Australia; ²Prince Henry's Institute of Medical Research, PO Box 5152 Clayton, Victoria 3168 Australia; ³MRC Centre for Reproductive Health, Queens Medical Research Institute, University of Edinburgh, Edinburgh, UK

Running title: Dynamics of the primordial follicle reserve

Corresponding author: Dr Jeff Kerr, Dept of Anatomy & Developmental Biology, Monash University, Clayton, Victoria 3800, Australia
P +613 99052723
E jeff.kerr@monash.edu

Abstract

The female germline comprises a reserve population of primordial (non-growing) follicles containing diplotene oocytes arrested in the first meiotic prophase. By convention the reserve is established when all individual oocytes are enclosed by granulosa cells. This commonly occurs prior to or around birth, according to species.

Histologically the “reserve” is the number of primordial follicles in the ovary at any given age and is ultimately depleted by degeneration and progression through folliculogenesis until exhausted. How and when the reserve reaches its peak number of follicles is determined by ovarian morphogenesis and germ cell dynamics involving i) oogonial proliferation and entry into meiosis producing an oversupply of oocytes, and ii) large-scale germ cell death resulting in markedly reduced numbers surviving as the primordial follicle reserve. Our understanding of the processes maintaining the reserve come primarily from genetically engineered mouse models, experimental activation or destruction of oocytes, and quantitative histological analysis. As the source of ovulated oocytes in postnatal life, the primordial follicle reserve requires regulation of i) its survival or maintenance, ii) suppression of development (dormancy) and iii) activation for growth and entry into folliculogenesis. The mechanisms influencing these alternate and complex inter-related phenomena remain to be fully elucidated. Drawing upon direct and indirect evidence, we discuss the controversial concept of postnatal oogenesis. This posits a rare population of oogonial stem cells that contribute new oocytes to partially compensate for the age-related decline in the primordial follicle reserve.

Introduction

The concept of a non-renewable primordial follicle pool, assembled around the time of birth in rodents and during gestation in humans, underpins a finite reproductive lifespan and is central to current understanding of ovarian biology. Consideration of the dynamics of the primordial follicle reserve raises more questions than there are answers, however, as although key pathways are emerging, their overall regulation and integration is poorly

understood. The main concepts include i) how the reserve is established, ii) processes causing elimination, iii) regulation of follicle-oocyte dormancy or activation into a growth phase, and iv) possibility of renewal accompanying the age-dependent decline. The significance of the dynamics of the reserve is no more apparent than during ovarian morphogenesis and germ cell development in prenatal life in humans, and perinatally in the mouse and rat. In these growth periods germ cells are produced in large numbers but many are subsequently eliminated, the outcome of which establishes the traditionally-defined primordial reserve. Mechanisms must exist to ensure that the majority of follicles are held intact and remain poised to participate in follicle growth, which in the human is preserved for decades. The reserve faces yet other challenges to its survival from exogenous agents that pose a risk of damage to the oocyte genome with accompanying DNA mutations, or more subtle epigenetic changes. How healthy or faulty oocytes within the reserve are recognized and respectively either preserved or destroyed is a key element impacting the dynamics of the primordial follicles. A reassessment of ‘topping up’ the reserve by the addition of new primordial follicles from ovarian germline stem cells has emerged in the past ten years. Although this concept has generated a lively debate and a resolution is far from complete, it introduces another factor that potentially affects the dynamics of the reserve. In summary, from the events that shape the establishment of the reserve in prenatal or neonatal ovaries up to the point of its functional exhaustion in adult life, we revisit the concepts of primordial follicle dynamics in the light of recent evidence influencing its stability, depletion or supplementation.

Establishing the primordial follicle reserve

The human fetal ovary

The developing fetal ovary supports the proliferation and maturation of germ cells and their development into primordial follicles. Studies by Block (Block 1951, Block 1952, Block 1953) of fetal, neonatal and adult human ovaries (n=53) using quantitative histological methods provided the first credible estimates of the number of primordial follicles. At 7-9 months gestation (n=10) he reported a range of 350,000 – 1.1 million primordial follicles per pair of ovaries, the average being about 700,000. In postnatal life from 6-9 years (n=5), the average was 500,000 declining to 8,000 between 40-44 years (n=7; (Block 1952)). The age-related fall in follicle supply was not discussed in Block's studies. Within a decade this oversight was corrected when in 1963 a landmark paper by Baker estimated the numbers of all germ cell types (normal and atretic) in human fetal ovaries (n=14). He calculated up to 6.8 million germ cells per pair of ovaries at 5 months gestation declining to about 2 million at the time of birth. The scale of germ cell loss was comparable to the germ cell attrition reported for the rat ovary (Holmes & Mandl 1962, Beaumont & Mandl 1963) suggesting common regulatory mechanisms governing the perinatal supply of primordial follicles. Recent analyses using more accurate stereological methods have expanded on the rate and extent of germ cell proliferation up to 19 weeks gestation (Mamsen *et al.* 2011), reaching nearly 5 million germ cells per ovary at that time, although no distinction is made between stages of development of the germ cells and the extent of inclusion within primordial follicles. Primordial follicles are formed from about 15 weeks gestation in the human fetal ovary (Fig. 1) based on the association of diplotene oocytes with pregranulosa cells (Baker 1963, Forabosco &

Sforza 2007). Their number steadily rises during the second trimester, and plateaus in the third trimester with approximately 350,000-400,000 per ovary at birth. From about 22 weeks, some primordial follicles activate to form the first growing or primary follicles (Maheshwari & Fowler 2008). As far as we know, **the second trimester period of human fetal ovarian development** is the only phase in the history of the dynamics of the reserve where it is increasing in overall number by the addition of oocytes surviving to reach diplotene arrest of meiosis I.

Little is known about the factors responsible for producing this excess of germ cells in the fetal ovary. Array-based studies have described the transcriptome in human fetal ovaries (Fowler *et al.* 2009), potentially allowing identification of regulatory pathways. A network of interacting oocyte transcription factors crucial for oocyte survival and development around the time of follicle formation has been described in the mouse using knock-out models (**Dong *et al.* 1996, Rajkovic *et al.* 2004, Pangas *et al.* 2006**), with some, such as FIGLA, demonstrated to have comparable expression in the human ovary (Huntriss *et al.* 2002, Bayne *et al.* 2004). Limited functional studies of human fetal ovaries have identified activin A (Martins da Silva *et al.* 2004, Coutts *et al.* 2008, Childs & Anderson 2009) and neurotrophin pathways (Anderson *et al.* 2002, Spears *et al.* 2003, Childs *et al.* 2010a) as likely key determinants of oogonial survival and proliferation and follicle formation (Fig.2). Activin β A is expressed by germ cells in nests and *in vitro* exposure to activin A promotes germ cell survival (Martins da Silva *et al.* 2004). Activin β A expression is lost immediately prior to nest breakdown and follicle formation (Coutts *et al.* 2008), and it is thought that this might act as switch allowing follicle formation

involving the de-repression of kit ligand expression (Childs & Anderson 2009). In the mouse, activin A administration *in utero* increased primordial follicle number after birth, although this difference was lost later in life (Bristol-Gould *et al.* 2006a). The BMPs have been suggested to positively regulate oogonial proliferation and survival in the mouse (Pesce *et al.* 2002), but in contrast experimental human data suggests that BMP4 increases germ cell apoptosis (Childs *et al.* 2010b), possibly explained by differences in experimental methodology (i.e. isolated germ cells in the mouse vs in their physiological niche in human whole ovary studies).

The neurotrophins BDNF and NT4 are expressed by ovarian somatic cells within the cell nests (i.e. presumed precursors to granulosa cells: Fig.2) with both ligands expressed in human but only NT4 in mouse. Mouse knock-out models of the TrkB receptor, targeted by both BDNF and NT4, have resulted in phenotypes including loss of oocytes at the time of follicle formation (Spears *et al.* 2003) and loss of initiation of follicle growth (Paredes *et al.* 2004). Oocyte-derived activin β A regulates BDNF expression in human ovarian somatic cells, and NT4 expression in mouse (Childs *et al.* 2010a), exemplifying a pathway by which the oocyte regulates the surrounding somatic environment, and also demonstrating a conserved pathway between species although involving diverse mediators. Prostaglandin E2 acting on oocytes may also contribute to the regulation of expression of activin β A and BDNF (Bayne *et al.* 2009), and there are undoubtedly other pathways involved. These interactions, derived from experimental human tissue studies, are illustrated in Figure 2.

More is known about the circumstances of oocyte death. We use the term ‘circumstances’ because of the limited opportunities available for analysis of human material (and no

prospects for *in vivo* experimentation) with most of our knowledge derived from the mouse. Although there are numerous descriptions of specific germ cell types and the timing of their demise in the human fetal ovary that impact on the dynamics of the reserve, the mechanisms responsible remain largely unknown (Maheshwari & Fowler 2008, Hartshorne *et al.* 2009). Much attention has focused on apoptosis (Vaskivuo *et al.* 2001, Fulton *et al.* 2005, Poljicanin *et al.* 2012), although emerging evidence also suggests that the mode of germ cell elimination, especially in meiosis, may be ovary-specific and occurs by several mechanisms not limited to the classic apoptotic pathways (Abir *et al.* 2002). Efforts to identify and quantitate the characteristics of apoptosis as a principal or coherent explanation for oocyte depletion in the human fetal ovary often demonstrate the difficulties and inconsistencies in interpretation of cause and effect, probably due to differential gene expression among cell populations that may be at rest, proliferating, maturing, dying or phagocytosing (Kurilo 1981, De Pol *et al.* 1997, Vaskivuo *et al.* 2001, Abir *et al.* 2002, Hartley *et al.* 2002, Fulton *et al.* 2005, Stoop *et al.* 2005, Albamonte *et al.* 2008, Jaaskelainen *et al.* 2010, Boumela *et al.* 2011, Poljicanin *et al.* 2012). Nevertheless, these and other studies demonstrate that the Bcl-2 gene family is an important regulator (among other factors) of the balance between survival or death of oocytes prior to primordial follicle formation.

The embryonic and neonatal mouse ovary

Germ cells of the embryonic mouse ovary follow a similar pattern of development as in the human except that it is only after birth that oocytes are fully assembled into the primordial follicle reserve, usually within 2-3 days (Fig 3). In common with the human fetal ovary there is a significant oversupply of oocytes entering meiosis prior to birth,

which is markedly reduced in the perinatal period of development (Fig 4; (Peters *et al.* 1978, McClellan *et al.* 2003, Kerr *et al.* 2006, Pepling 2006, Pepling *et al.* 2010). With the advantage of experimental interventions such as the ability to modify gene expression, **much** of our knowledge regarding female germ cell death mechanisms has been generated in the mouse.

Because primordial follicle formation is associated with significant germ cell attrition (Kezele *et al.* 2002, Pepling 2006), investigations into the associated death mechanisms have been topical and numerous laboratories, using both *in vivo* and *in vitro* techniques have concluded that apoptosis (Coucouvani *et al.* 1993, De Pol *et al.* 1997, Pepling & Spradling 2001, De Felici *et al.* 2008, Xu *et al.* 2011) autophagy (Lobascio *et al.* 2007, De Felici *et al.* 2008, Rodrigues *et al.* 2009), and direct extrusion from the ovaries (Rodrigues *et al.* 2009) are all contributory mechanisms of pre- and neonatal oocyte demise. Apoptosis, the most favoured of the three, has been demonstrated not only in mouse models directly targeting Bcl-2 and caspase genes (Bergeron *et al.* 1998, Perez *et al.* 1999, Rucker *et al.* 2000, Flaws *et al.* 2001, Flaws *et al.* 2006, Alton & Taketo 2007, Ghafari *et al.* 2007, Greenfeld *et al.* 2007, Gursoy *et al.* 2008, Ghafari *et al.* 2009) but also because of the findings from several gene knockout (or overexpressor) models belonging to the TNF pathway (Marcinkiewicz *et al.* 2002, Greenfeld *et al.* 2007), PAR family (Wen *et al.* 2009), and TGF β family (Kimura *et al.* 2011), all of which actively participate in oocyte loss by regulating apoptosis.

What controls oocyte death to establish the reserve?

185 For oogonia and oocytes the mechanism of cell death implemented may be related to the
 186 signal to die. Most studies of oocyte dynamics in the neonatal mouse ovary point to
 187 apoptosis as the mode of death (Ghafari *et al.* 2009, Boumela *et al.* 2011, Hu *et al.* 2011.)
 188 Therefore, the primordial follicle reserve is presumably established by a balance between
 189 the availability of a large number of germ cells and subsequent programmed cell death.
 190 Why so many oocytes are produced only to be eliminated remains a mystery, but some
 191 possibilities are i) failure of mitosis/meiosis involving **defective** chromosome spindle
 192 **functions**, ii) unrepaired DNA damage, iii) insufficient pregranulosa cells, and iv)
 193 **degeneration of oocytes during** restructuring of oocyte **cysts** or nests into primordial
 194 follicles. The first clues that one member of the p53 gene network had a significant role
 195 in controlling oocyte fate came from studies showing that p63, specifically the TAp63 α
 196 isoform, is expressed uniquely in mouse oocytes and is responsible for their elimination if
 197 for example their DNA is damaged (Suh *et al.* 2006). Thus p63 has a role in regulating
 198 oocyte survival to establish the primordial follicle reserve. Its expression in late prophase
 199 I oocytes but not in early meiotic oocytes or oogonia in fetal ovaries (both mouse and
 200 human), suggests a universal role for p63 in protection of the female germline
 201 represented by the primordial reserve (Livera *et al.* 2008). In the early postnatal mouse
 202 ovary p63 controls oocyte supply by transcriptional induction of BH3-only proteins
 203 PUMA or PUMA and NOXA combined (Kerr *et al.* 2012b). These pro-apoptotic Bcl-2
 204 members can initiate oocyte apoptosis either by direct or indirect activation of BAX and
 205 BAK. Deletion of *Puma* or *Puma* and *Noxa* together results in an oversupply of
 206 primordial follicles in postnatal day 10 mouse ovaries, and deletion of other BH3-only
 207 genes, *Bmf* or *Bim* also amplifies the reserve with up to triple the numbers of oocytes

compared with age-matched controls (Fig. 5). The role if any of the other BH3-only proteins remains unknown. Given that ‘overstocking’ of the primordial reserve in the mouse ovary is wholly or partly the net result of a balance between pro- and anti-apoptotic events, it remains to be shown at what time and which germ cell types (i.e. oogonia and/or oocytes) are affected.

While these studies confirm that apoptotic regulatory mechanisms are key factors in altering the dynamics of the primordial reserve, they do not exclude the possibility of alternate or complementary processes for adjusting the oocyte population. Other studies of the developing human or mouse ovary have demonstrated that the apoptotic paradigm does not satisfactorily account for all aspects of germ cell death (Vaskivuo *et al.* 2001, Abir *et al.* 2002, Alton & Taketo 2007, De Felici *et al.* 2008, Rodrigues *et al.* 2009, Gawriluk *et al.* 2011). Alternative modes of cell death that may participate in oogonial-oocyte elimination include autophagy (Guillon-Munos *et al.* 2006, Rubinstein & Kimchi 2012), mitotic arrest (Wartenberg *et al.* 2001) or necroptosis (Vandenabeele *et al.* 2010, Christofferson & Yuan 2010).

Dynamics of the postnatal primordial follicle reserve and consequences for reproductive lifespan

Analogous to a stockpile of a precious resource, most oocytes of the primordial reserve are retained as quiescent follicles to support future ovulations throughout the reproductive lifespan. A poorly stocked initial reserve or one in which primordial

231 follicles are precociously depleted, will result in infertility and in the human, a shortened
232 reproductive lifespan and early menopause (Nelson *et al.* 2013). Current concepts involve
233 progressive loss of human female fertility expressed through subfertility, sterility and the
234 menopause at approximately 10 year intervals (Broekmans *et al.* 2009) . Thus a
235 menopause at age 40 (the traditional definition of the upper limit of ‘premature’) implies
236 a loss of fertility at 30 and falling fertility from the early 20s. Mathematical analyses of
237 the age-related decline of the non-growing follicle (NGF) reserve (ie. primordial follicles)
238 in human ovaries predicts that if at birth one ovary had 35,000 NGFs, menopause would
239 occur at around 40 years of age but would be delayed to 60 years if the ovary began with
240 2.5million NGFs (Wallace & Kelsey 2010, Kelsey *et al.* 2012). The number and types of
241 molecules believed to maintain the balance between quiescence and activation of the
242 primordial follicle reserve continue to be discovered chiefly from the study of transgenic
243 mouse models (Reddy *et al.* 2010, Kim 2012, Monget *et al.* 2012, Pangas 2012, Adhikari
244 *et al.* 2013). A key pathway implicated in this is the PI3K pathway, which may have a
245 crucial integrative role linking many of the factors associated with the balance between
246 follicle growth suppression, activation, and the maintenance of healthy quiescence (Fig.
247 6). Molecules in this pathway include the tuberous sclerosis complex 1 (TSC1) which
248 interacts with phosphatase and tensin homolog deleted on chromosome 10 (PTEN) to
249 maintain quiescence, and the mammalian target of rapamycin (mTORC) which is an
250 activator, and negatively regulated by TSC1 (Zheng *et al.* 2012). Both the oocyte and its
251 pre-granulosa cells are the source and probably the targets for these factors that
252 physiologically exert both stimulatory and inhibitory actions upon the primordial follicle
253 reserve. In addition to intracrine (**factors produced and acting within a cell**) and/or

paracrine inhibition of the recruitment of primordial follicles, an additional ‘brake’
 maintaining their quiescence and perhaps regulating the rate of recruitment may be
 applied by the growing follicle pool (Barnett *et al.* 2006, Moniruzzaman & Miyano 2010,
 Reddy *et al.* 2010, Monget *et al.* 2012). Mathematical modeling of histomorphometric
 data has shown age-dependent differential rates of NGF recruitment in the postnatal
 human ovary (Wallace & Kelsey 2010) with the great majority of follicles lost in the
 younger years. Implicit for these observations is the concept that in the early phases of
 postnatal life including and beyond puberty, some intra-ovarian mechanism limits the
 decline of the primordial reserve to conserve its stockpile of follicles. In the postnatal
 mouse ovary it has been suggested that the preservation of a set range of follicle number
 in the primordial reserve is consistent with a ‘quorum-sensing’ model (Bristol-Gould *et al.*
 2006b, Tingen *et al.* 2009). In this model the ovary can eliminate excess primordial
 follicles perhaps via a Bcl-2 cell death mechanism but on current evidence cannot add
 primordial follicles to an otherwise abnormally insufficient reserve. **While biochemical**
pathways that seem to be involved in the maintenance of primordial follicle health
have been proposed based on knock-out models (eg Pdk1 and Rps6: (Reddy *et al.*
2009), how (or indeed whether) primordial follicle health is monitored
physiologically is an important but unclear question.
 What is the evidence for a ‘brake’ applied (at least temporarily) to the disappearance, by
 growth initiation or direct atresia, of primordial follicles from the reserve? In the Bl/6
 mouse strain, following the precipitous decline during the neonatal period, the depletion
 of primordial follicles per ovary is minimal, losing on average less than 1 follicle per day
 for up to 14 weeks (Kerr *et al.* 2006, Rodrigues *et al.* 2009) but thereafter declines

277 significantly up to 300 days (Kerr *et al.* 2012a). **Using cell lineage-tracing Lei &**
 278 **Spradling {Lei, 2013 #1764} showed that the primordial follicle population is highly**
 279 **stable in the postnatal mouse ovary. With an estimated half-life of 10 months in**
 280 **adult life, the supply of primordial follicles established in the neonatal ovary is**
 281 **sufficient to sustain adult folliculogenesis (and fertility) without a source of renewal**
 282 **({Lei, 2013 #1764}).** When growth-initiated i.e. primary follicles are counted, these
 283 decline significantly losing about 2.5 follicles on average per day (unpublished data).
 284 Could the growing primary follicles and their successors the secondary/antral follicles
 285 play a role in restraining recruitment from the primordial reserve? The preferential
 286 location of the reserve to the ovarian cortex with growing follicles mostly confined to the
 287 medulla (Da Silva-Buttkus *et al.* 2009) suggests a follicle-derived gradient of inhibitory
 288 and stimulatory signals that reflects this arrangement. Spatial analysis of primordial
 289 follicles has led to the proposal that these follicles inhibit each other by producing as yet
 290 unidentified paracrine factors that prevent their activation into primary follicles (Da
 291 Silva-Buttkus *et al.* 2009). Perhaps growing follicles influence the rate of entry of
 292 primordial follicles into the growth phase, and the phenotype of the AMH knock-out
 293 mouse suggests that AMH may contribute to this (Durlinger *et al.* 1999). Analysis of
 294 AMH concentrations in relation to NGF number and recruitment across life indicate
 295 changing relationships during puberty and early adult life (Fleming *et al.* 2012) in
 296 keeping with this factor also playing a significant role in the human. The signal for
 297 activation of a reserve follicle may also be based on the origin of the pregranulosa cells
 298 and timing of follicle formation, with a separate medullary population formed
 299 immediately after birth distinct from the cortical population that supports adult fertility

(Mork *et al.* 2012). This interpretation, based on mouse experimental data, appears to differ from a recent reanalysis of bovine ovarian development (Hummitzsch *et al.* 2013), which indicates that all pregranulosa cells arise early from precursor cells first identifiable within the ovarian surface epithelium.

Thus in the mouse, particularly during the early phase of reproductive life, oocytes destined for ovulation may in theory be supplied mainly from the diminishing primary follicle population. As time passes this temporary stock of growing follicles can by itself no longer sustain the folliculogenic production line and the dwindling size of the early growing follicle population becomes insufficient to exert an inhibitory affect or restraint over the primordial reserve. At that point some of the previously dormant primordial follicles are activated, and the reserve is mobilized. Accessing primordial follicles stored in the reserve will lead ultimately to its depletion whereupon folliculogenesis is curtailed and ovulation ceases. Such detailed information is not available from human studies, which can only be based on cross-sectional analysis of limited data sets. While an increase in the rate of follicle depletion with age is often cited and holds true when expressed as a proportion of remaining follicles, a recent mathematical analysis of the number of follicles leaving the non-growing pool shows that this increases through childhood, peaking at approximately 900 follicles per month at age 14 (with an average follicle endowment), then falling to 600 per month at age 25 and 200 per month at age 35 (Kelsey *et al.* 2012).

The primordial follicle reserve: is it renewable?

323 In 2004 Johnson et al proposed that in the mouse ovary, the incidence of ongoing, age-
324 related follicle elimination by atresia outstripped the contemporaneous supply available
325 in the primordial follicle reserve. This imbalance was predictive of exhaustion of the
326 reserve within a few weeks beyond puberty (Johnson *et al.* 2004), yet mice may remain
327 fertile for up to 12 months (Gosden *et al.* 1983). To offset the proposed loss of primordial
328 follicles evidence was presented for the existence of ovarian germline stem cells (GSC)
329 capable of proliferation and meiotic maturation into newly-minted oocytes (Johnson *et al.*
330 2004). Candidate cells were identified in the ovarian surface epithelium leading to the
331 opinion that GSC had been discovered in the mouse (Spradling 2004). Later the notion
332 that GSC arise from the surface epithelium was **revised because the small number**
333 **(6±3) estimated to be present in the postnatal day 40 ovary was insufficient to**
334 **generate new oocytes to offset normal follicle loss** (Johnson *et al.* 2005). Other studies
335 of the superficial ovarian cortex reported a mixed population of oocytes, primordial
336 follicles, oogonial-type cells and unidentified cells in mitosis (Kerr *et al.* 2006). In
337 **seeking an alternative source of GSC external to the ovaries**, an origin from bone
338 marrow and blood was next proposed with GSC seeding the mouse ovary to replenish the
339 natural decline in the primordial reserve oocytes (Johnson *et al.* 2005). **This study also**
340 **reported that in ovaries of mice** exposed to the cytotoxins doxorubicin (DXR) or
341 histone deacetylase inhibitor trichostatin A (TSA), resulted within 24-36hrs in respective
342 ‘spontaneous regeneration’ of lost primordial follicles or doubling of their numbers by
343 ‘de novo oocyte production’. Together these results were said to reinforce the concept
344 that oogenesis and folliculogenesis could occur in the adult ovary (Johnson *et al.* 2005).
345 **However other studies of the effects of DXR or TSA on mouse ovaries have shown**

depletion of the primordial follicle reserve with no evidence for regeneration (Kujjo *et al.* 2011, Kerr *et al.* 2012a). The contrasting outcomes of gain or loss of primordial follicles reported in different studies adds to the debate on the renewability of germ cells/oocytes in the postnatal ovary, and it remains the case that even if there is some physiological follicular renewal it too is finite (the incontrovertible existence of the menopause), whether as a result of limiting supply of germ cells, required associated somatic cells or both. A parabiosis model (Eggan *et al.* 2006) did not provide supportive evidence for a bone marrow or blood-borne source for ovulated mouse oocytes, but the presence or absence in the ovaries, of marrow- or blood-derived GSC or new follicles was not investigated. When bone marrow obtained from transgenic mice expressing germline-specific green fluorescent protein (GFP) was transplanted into wild-type recipients, GFP-positive germ cells/oocytes were detected in recipient ovaries albeit at a low frequency of $1.4 \pm 0.6\%$ of the total immature follicle pool but none developed into ovulated oocytes (Lee *et al.* 2007). Further studies of the identification and developmental potential of GSC or oogonial stem cells (OSC) in the mouse and human ovary are now available (Zou *et al.* 2009, Pacchiarotti *et al.* 2010, White *et al.* 2012, Zhang *et al.* 2012) but the interpretation of the results continues to generate controversy (Oatley & Hunt 2012, Woods *et al.* 2013). The human data thus far available (White *et al.* 2012) indicate the existence of a small number of cells within the ovary that can be extracted, proliferate *in vitro*, and after labeling and injection into isolated human ovarian cortex tissue, formed primordial follicles containing labeled oocytes. In the mouse, injection of these cells into adult ovary resulted in the ovulation of fertilizable oocytes and livebirths (Zou *et al.*

2009, White *et al.* 2012). This work requires further corroboration, and while potentially of considerable scientific and medical interest, provides no evidence that these cells contribute to physiological ovarian function, including fertility.

If OSC conform to stem cell kinetics they must proliferate by mitosis to preserve their 'stemness'. Genomic analysis in mice of the number of preceding mitotic divisions for antral follicle oocytes revealed how many germ cell divisions have occurred since the zygote stage, this being referred to as oocyte 'depth' (Reizel *et al.* 2012). This study found that oocyte depth increases with age; 13 divisions on average in oocytes sampled at day 30 but 20 divisions in oocytes obtained at 350 days. Do these divisions occur only during embryonic development or throughout all of life?

The first possibility is consistent with the 'production-line' hypothesis (Henderson & Edwards 1968) i.e. the order in which oocytes ovulate postnatally follows the order in which oogonia entered meiosis (and cannot re-enter mitosis) in the embryonic ovary. Meiotic entry is not an 'all-or-none' event but a gradual process occurring from e13.5-e18.5 (Peters *et al.* 1962, Ghafari *et al.* 2007, Ghafari *et al.* 2009) and progressing in the ovary in a cranial-caudal direction (Bullejos & Koopman 2004). Many oogonia in the fetal mouse (and human) ovary continue mitosis whilst others enter meiotic prophase (Evans 1982, Fulton *et al.* 2005) and therefore oogonia with fewer or greater numbers of mitotic divisions would respectively transition early or later into meiosis. Medullary oocytes become early-activated primordial follicles but cortex-resident oocytes are delayed in their assembly as primordial follicles (Fig. 3B). This pattern of germ cell distribution and subsequent dynamics is initiated in the mouse ovary at e13.5 (Byskov *et al.* 1997).

(Woods *et al.* 2012) favour the alternative possibility whereby additional mitoses of OSC during postnatal life produce oocytes of greater ‘depth’ consistent with measured genetic signatures. Cells with OSC-type properties have been found among primordial follicles in or subjacent to the surface epithelium of the neonatal mouse ovary (Zou *et al.* 2009) and although cells with similar characteristics have been observed (Kerr *et al.* 2006) their identity, function and fate remain to be confirmed. Bristol-Gould *et al.* (Bristol-Gould *et al.* 2006a) and Tingen *et al.* (Tingen *et al.* 2009) reported that 5% of germ cells in the neonatal mouse ovary are ‘residual’ oogonia, which did not enter meiosis between e13.5-e18.5. If bypassing oocyte nest formation and encapsulation to form primordial follicles, do these orphan oogonia represent the OSC, being rare, unrecognized with routine histology (not being primordial follicles) and problematic to characterize using established stem cell or germline cell markers? Further investigations may reveal if these reputed OSC co-exist with the conventional primordial follicle reserve and represent a hitherto unknown population of germ cells with the potential of development given special opportunity.

Conclusions

From the time of its formation and development within the fetal or neonatal ovary, and throughout the postnatal reproductive lifespan, the primordial follicle reserve is subject to constant change. The remarkable increase then substantial loss of germ cells in the fetal ovary impacts the dynamics of the reserve to the extent of providing oocytes for assembly into primordial follicles. The maximum supply of primordial follicles is the net result of the addition to the reserve of suitably developed oocytes, counterbalanced by depletion

through germ cell death, and depending on species, activation of follicles into a growth phase. Mechanisms controlling germ cell proliferation are not fully understood but evidence is emerging for regulation by interactions between a variety of transcription and growth factors. Elimination of germ cells is likely due to several processes particularly via apoptosis but with increasing evidence for non-apoptotic cell death, such as autophagy, acting alone or in combination with apoptosis and dependant on the type and biological status of the germ cells necessitating their removal. Although in postnatal life many primordial follicles in humans may be preserved for decades in a state of dormancy, the dynamic nature of the primordial follicle reserve is again evident, chiefly through depletion as follicles activate and enter folliculogenesis, and possibly by direct elimination/atresia of those follicles sustaining genomic impairment. **Theoretically, manipulation of the rate of activation of primordial follicle pool could be of clinical value. Temporary increased activation could be of value to women requiring assisted conception later in life to increase the number of oocytes that could be recovered, and conversely slowed activation could be of value to delay the menopause and possibly prolong natural fertility if a reduced pool (and hence increased risk of early menopause) was identified. These possibilities remain remote and, as with all manipulations of the germ line, raise very serious safety considerations.** Recent reports of the existence of a rare population of germline stem cells in mouse and human ovaries have led to suggestions that these cells may partially replenish the reserve as its primordial follicle supply is diminished. If further work confirms recent studies showing that isolated GSC can form follicles with fertilizable

oocytes and viable embryos, this may usher in a new paradigm: an ancillary germ cell population coexisting with the primordial follicle pool, the ‘reserve’ of the reserve.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Acknowledgments

The authors work is supported by grants from the UK Medical Research Council (G1100357 to RAA) and a TM Ramsay Fellowship and Ramaciotti Establishment Grant (to MM).

Figure 1. Estimates of germ cell populations of the human fetal ovary based on histomorphometric analysis reported by Baker (1963) and Forabosco & Sforza (2007). Germ cells are always dying, the numbers of atretic germ cells (which include all oogonia and oocytes) being equal to or exceeding the numbers of individual diplotene oocytes or those forming primordial follicles. Primordial follicles begin to form at 15 weeks gestation and at birth the fetal ovary on average contains approximately 400,000 primordial follicles. This represents only 12% of the total germ cell number (healthy and atretic) present at 22 weeks gestation.

Figure 2. Human ovarian development and primordial follicle formation. Illustrative immunohistochemical images to demonstrate (A) proliferative oogonia (mitotic cells

identified by arrows) and oocytes (Oo) within germ cell nests with intermixed NT4-expressing (brown) pregranulosa cells, with the somatic cells (sc) of the stromal regions not expressing NT4 13 weeks gestation. (B) NT4 expression (brown) is also confined to pregranulosa cells within oocyte (Oo) nests and the granulosa cells of newly-formed primordial follicles, with no expression in stromal cells (sc): 21 weeks gestation. (C) Activin βA is expressed by some nests of oocytes (Oo) (green; red nuclear counterstain) but with much weaker expression in others (arrow), indicating synchronous development of oocytes within a nest; 19 weeks gestation. (D) Schematic representation of experimentally-derived interactions between growth factors expressed by oogonia/oocytes of the human fetal ovary and the adjacent pregranulosa cells. Stimulatory (+) and inhibitory (-) regulation as indicated. Scale bar A-C, 20 μ m.

Figure 3. Development of oocytes and primordial follicles. (A) Pachytene oocytes in e17 mouse ovary showing their arrangement into nests in which individual oocytes are not enclosed by somatic cells that will later become pregranulosa cells of primordial follicles. Scale bar 15 μ m. (B) Postnatal day 1 mouse ovary showing oocyte nests in the cortex region to the right, and larger individual primordial follicles (pf) in the medulla. Pyknotic structures (example at arrowhead) represent degenerative oocytes. Scale bar 20 μ m.

Figure 4. Schematic diagram illustrating the general trends of endowment of oocytes and primordial follicles based on stereological analysis in the Bl/6 mouse ovary over indicated ages. Dashed line: is not data based but an estimation of germ cell increase; solid line: mostly based on published data. Oocyte number increases markedly towards

the end of fetal life but many are lost as they assemble to form primordial follicles in the first days after birth. For up to two weeks postnatally primordial follicles decline significantly then enter a period of very slow follicle loss for up to several months followed again by renewed depletion until near or total exhaustion around 12 months of age. Data based in part on Myers *et al.* 2004, Kerr *et al.* 2006, 2012a, Rodrigues *et al.* 2009, Lei & Spradling 2013.

Figure 5. Gene regulation of the primordial follicle reserve in the mouse ovary. Comparison of primordial follicle supply in postnatal day 10 mouse ovaries in wild-type, *p53* ^{-/-} and various BH3 member knockout models. Based on data from Kerr *et al* 2012b.

Figure 6. Schematic diagram illustrating the options available for primordial follicles leaving the arrested reserve (growth/suppression of growth; maintenance of health; elimination and possibly renewal) and representative proteins or genes identified as regulators of these various pathways.

References

- Abir R, Orvieto R, Dicker D, Zukerman Z, Barnett M & Fisch B** 2002 Preliminary studies on apoptosis in human fetal ovaries. *Fertil Steril* **78** 259-264.
- Adhikari D, Risal S, Liu K & Shen Y** 2013 Pharmacological inhibition of mTORC1 prevents over-activation of the primordial follicle pool in response to elevated PI3K signaling. *PLoS One* **8** e53810.
- Albamonte MS, Willis MA, Albamonte MI, Jensen F, Espinosa MB & Vitullo AD** 2008 The developing human ovary: immunohistochemical analysis of germ-cell-specific VASA protein, BCL-2/BAX expression balance and apoptosis. *Hum Reprod* **23** 1895-1901.

- 507 **Alton M & Taketo T** 2007 Switch from BAX-dependent to BAX-independent germ cell
508 loss during the development of fetal mouse ovaries. *J Cell Sci* **120** 417-424.
- 509 **Anderson RA, Cambray N, Hartley PS & McNeilly AS** 2002 Expression and
510 localization of inhibin alpha, inhibin/activin betaA and betaB and the activin type
511 II and inhibin beta-glycan receptors in the developing human testis. *Reproduction*
512 **123** 779-788.
- 513 **Baker TG** 1963 A Quantitative and Cytological Study of Germ Cells in Human Ovaries.
514 *Proc R Soc Lond B Biol Sci* **158** 417-433.
- 515 **Barnett KR, Schilling C, Greenfeld CR, Tomic D & Flaws JA** 2006 Ovarian follicle
516 development and transgenic mouse models. *Hum Reprod Update* **12** 537-555.
- 517 **Bayne RA, Eddie SL, Collins CS, Childs AJ, Jabbour HN & Anderson RA** 2009
518 Prostaglandin E2 as a regulator of germ cells during ovarian development. *J Clin*
519 *Endocrinol Metab* **94** 4053-4060.
- 520 **Bayne RA, Martins da Silva SJ & Anderson RA** 2004 Increased expression of the
521 FIGLA transcription factor is associated with primordial follicle formation in the
522 human fetal ovary. *Mol Hum Reprod* **10** 373-381.
- 523 **Beaumont HM & Mandl AM** 1963 A Quantitative Study of Primordial Germ Cells in
524 the Male Rat. *J Embryol Exp Morphol* **11** 715-740.
- 525 **Bergeron JM, Gahr M, Horan K, Wibbels T & Crews D** 1998 Cloning and in situ
526 hybridization analysis of estrogen receptor in the developing gonad of the red-
527 eared slider turtle, a species with temperature-dependent sex determination. *Dev*
528 *Growth Differ* **40** 243-254.
- 529 **Block E** 1951 Quantitative morphological investigations of the follicular system in
530 women; methods of quantitative determinations. *Acta Anat (Basel)* **12** 267-285.
- 531 **Block E** 1952 Quantitative morphological investigations of the follicular system in
532 women; variations at different ages. *Acta Anat (Basel)* **14** 108-123.
- 533 **Block E** 1953 A quantitative morphological investigation of the follicular system in
534 newborn female infants. *Acta Anat (Basel)* **17** 201-206.
- 535 **Boumela I, Assou S, Aouacheria A, Haouzi D, Dechaud H, De Vos J, Handyside A &**
536 **Hamamah S** 2011 Involvement of BCL2 family members in the regulation of
537 human oocyte and early embryo survival and death: gene expression and beyond.
538 *Reproduction* **141** 549-561.
- 539 **Bristol-Gould SK, Kreeger PK, Selkirk CG, Kilen SM, Cook RW, Kipp JL, Shea**
540 **LD, Mayo KE & Woodruff TK** 2006a Postnatal regulation of germ cells by
541 activin: the establishment of the initial follicle pool. *Dev Biol* **298** 132-148.
- 542 **Bristol-Gould SK, Kreeger PK, Selkirk CG, Kilen SM, Mayo KE, Shea LD &**
543 **Woodruff TK** 2006b Fate of the initial follicle pool: empirical and mathematical
544 evidence supporting its sufficiency for adult fertility. *Dev Biol* **298** 149-154.
- 545 **Broekmans FJ, Soules MR & Fauser BC** 2009 Ovarian aging: mechanisms and clinical
546 consequences. *Endocr Rev* **30** 465-493.
- 547 **Bullejos M & Koopman P** 2004 Germ cells enter meiosis in a rostro-caudal wave during
548 development of the mouse ovary. *Mol Reprod Dev* **68** 422-428.
- 549 **Byskov AG, Guoliang X & Andersen CY** 1997 The cortex-medulla oocyte growth
550 pattern is organized during fetal life: an in-vitro study of the mouse ovary. *Mol*
551 *Hum Reprod* **3** 795-800.

- 552 **Childs AJ & Anderson RA** 2009 Activin A selectively represses expression of the
 553 membrane-bound isoform of Kit ligand in human fetal ovary. *Fertil Steril* **92**
 554 1416-1419.
- 555 **Childs AJ, Bayne RA, Murray AA, Martins Da Silva SJ, Collins CS, Spears N &**
 556 **Anderson RA** 2010a Differential expression and regulation by activin of the
 557 neurotrophins BDNF and NT4 during human and mouse ovarian development.
 558 *Dev Dyn* **239** 1211-1219.
- 559 **Childs AJ, Kinnell HL, Collins CS, Hogg K, Bayne RA, Green SJ, McNeilly AS &**
 560 **Anderson RA** 2010b BMP signaling in the human fetal ovary is developmentally
 561 regulated and promotes primordial germ cell apoptosis. *Stem Cells* **28** 1368-1378.
- 562 **Christofferson DE & Yuan J** 2010 Necroptosis as an alternative form of programmed
 563 cell death. *Curr Opin Cell Biol* **22** 263-268.
- 564 **Coucovanis EC, Sherwood SW, Carswell-Crumpton C, Spack EG & Jones PP**
 565 1993 Evidence that the mechanism of prenatal germ cell death in the mouse is
 566 apoptosis. *Exp Cell Res* **209** 238-247.
- 567 **Coutts SM, Childs AJ, Fulton N, Collins C, Bayne RA, McNeilly AS & Anderson**
 568 **RA** 2008 Activin signals via SMAD2/3 between germ and somatic cells in the
 569 human fetal ovary and regulates kit ligand expression. *Dev Biol* **314** 189-199.
- 570 **Da Silva-Buttkus P, Marcelli G, Franks S, Stark J & Hardy K** 2009 Inferring
 571 biological mechanisms from spatial analysis: prediction of a local inhibitor in the
 572 ovary. *Proc Natl Acad Sci U S A* **106** 456-461.
- 573 **De Felici M, Lobascio AM & Klinger FG** 2008 Cell death in fetal oocytes: many
 574 players for multiple pathways. *Autophagy* **4** 240-242.
- 575 **De Pol A, Vaccina F, Forabosco A, Cavazzuti E & Marzona L** 1997 Apoptosis of
 576 germ cells during human prenatal oogenesis. *Hum Reprod* **12** 2235-2241.
- 577 **Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N & Matzuk MM** 1996 Growth
 578 differentiation factor-9 is required during early ovarian folliculogenesis. *Nature*
 579 **383** 531-535.
- 580 **Durlinger AL, Kramer P, Karels B, de Jong FH, Uilenbroek JT, Grootegoed JA &**
 581 **Themmen AP** 1999 Control of primordial follicle recruitment by anti-Mullerian
 582 hormone in the mouse ovary. *Endocrinology* **140** 5789-5796.
- 583 **Eggan K, Jurga S, Gosden R, Min IM & Wagers AJ** 2006 Ovulated oocytes in adult
 584 mice derive from non-circulating germ cells. *Nature* **441** 1109-1114.
- 585 **Evans MA** 1982 Effect of phenytoin on calcium disposition in pregnant and nonpregnant
 586 mice. *Toxicol Appl Pharmacol* **63** 422-428.
- 587 **Flaws JA, Hirshfield AN, Hewitt JA, Babus JK & Furth PA** 2001 Effect of bcl-2 on
 588 the primordial follicle endowment in the mouse ovary. *Biol Reprod* **64** 1153-
 589 1159.
- 590 **Flaws JA, Marion SL, Miller KP, Christian PJ, Babus JK & Hoyer PB** 2006 Effect
 591 of bcl-2 overexpression in mice on ovotoxicity caused by 4-vinylcyclohexene.
 592 *Toxicol Appl Pharmacol* **215** 51-56.
- 593 **Fleming R, Kelsey TW, Anderson RA, Wallace WH & Nelson SM** 2012 Interpreting
 594 human follicular recruitment and antimullerian hormone concentrations
 595 throughout life. *Fertil Steril* **98** 1097-1102.
- 596 **Forabosco A & Sforza C** 2007 Establishment of ovarian reserve: a quantitative
 597 morphometric study of the developing human ovary. *Fertil Steril* **88** 675-683.

- Fowler PA, Flannigan S, Mathers A, Gillanders K, Lea RG, Wood MJ, Maheshwari A, Bhattacharya S, Collie-Duguid ES, Baker PJ, Monteiro A & O'Shaughnessy PJ** 2009 Gene expression analysis of human fetal ovarian primordial follicle formation. *J Clin Endocrinol Metab* **94** 1427-1435.
- Fulton N, Martins da Silva SJ, Bayne RA & Anderson RA** 2005 Germ cell proliferation and apoptosis in the developing human ovary. *J Clin Endocrinol Metab* **90** 4664-4670.
- Gawriluk TR, Hale AN, Flaws JA, Dillon CP, Green DR & Rucker EB, 3rd** 2011 Autophagy is a cell survival program for female germ cells in the murine ovary. *Reproduction* **141** 759-765.
- Ghafari F, Gutierrez CG & Hartshorne GM** 2007 Apoptosis in mouse fetal and neonatal oocytes during meiotic prophase one. *BMC Dev Biol* **7** 87.
- Ghafari F, Pelengaris S, Walters E & Hartshorne GM** 2009 Influence of p53 and genetic background on prenatal oogenesis and oocyte attrition in mice. *Hum Reprod* **24** 1460-1472.
- Gosden RG, Laing SC, Felicio LS, Nelson JF & Finch CE** 1983 Imminent oocyte exhaustion and reduced follicular recruitment mark the transition to acyclicity in aging C57BL/6J mice. *Biol Reprod* **28** 255-260.
- Greenfeld CR, Roby KF, Pepling ME, Babus JK, Terranova PF & Flaws JA** 2007 Tumor Necrosis Factor (TNF) Receptor Type 2 Is an Important Mediator of TNF alpha Function in the Mouse Ovary. *Biol Reprod* **76** 224-231.
- Guillon-Munos A, van Bemmelen MX & Clarke PG** 2006 Autophagy can be a killer even in apoptosis-competent cells. *Autophagy* **2** 140-142.
- Gursoy E, Ergin K, Basaloglu H, Koca Y & Seyrek K** 2008 Expression and localisation of Bcl-2 and Bax proteins in developing rat ovary. *Res Vet Sci* **84** 56-61.
- Hartley PS, Bayne RA, Robinson LL, Fulton N & Anderson RA** 2002 Developmental changes in expression of myeloid cell leukemia-1 in human germ cells during oogenesis and early folliculogenesis. *J Clin Endocrinol Metab* **87** 3417-3427.
- Hartshorne GM, Lyrakou S, Hamoda H, Oloto E & Ghafari F** 2009 Oogenesis and cell death in human prenatal ovaries: what are the criteria for oocyte selection? *Mol Hum Reprod* **15** 805-819.
- Henderson SA & Edwards RG** 1968 Chiasma frequency and maternal age in mammals. *Nature* **218** 22-28.
- Holmes RL & Mandl AM** 1962 The effect of norethynodrel on the ovaries and pituitary gland of adult female rats. *J Endocrinol* **24** 497-515.
- Hu W, Zheng T & Wang J** 2011 Regulation of Fertility by the p53 Family Members. *Genes Cancer* **2** 420-430.
- Hummitzsch K, Irving-Rodgers HF, Hatzirodos N, Bonner W, Sabatier L, Reinhardt DP, Sado Y, Ninomiya Y, Wilhelm D & Rodgers RJ** 2013 A new model of development of the mammalian ovary and follicles. *PLoS One* **8** e55578.
- Huntriss J, Gosden R, Hinkins M, Oliver B, Miller D, Rutherford AJ & Picton HM** 2002 Isolation, characterization and expression of the human Factor In the Germline alpha (FIGLA) gene in ovarian follicles and oocytes. *Mol Hum Reprod* **8** 1087-1095.

- 643 **Jaaskelainen M, Nieminen A, Pokkyla RM, Kauppinen M, Liakka A, Heikinheimo**
644 **M, Vaskivuo TE, Klefstrom J & Tapanainen JS** 2010 Regulation of cell death
645 in human fetal and adult ovaries--role of Bok and Bcl-X(L). *Mol Cell Endocrinol*
646 **330** 17-24.
- 647 **Johnson J, Bagley J, Skaznik-Wikiel M, Lee HJ, Adams GB, Niikura Y, Tschudy**
648 **KS, Tilly JC, Cortes ML, Forkert R, Spitzer T, Iacomini J, Scadden DT &**
649 **Tilly JL** 2005 Oocyte generation in adult mammalian ovaries by putative germ
650 cells in bone marrow and peripheral blood. *Cell* **122** 303-315.
- 651 **Johnson J, Canning J, Kaneko T, Pru JK & Tilly JL** 2004 Germline stem cells and
652 follicular renewal in the postnatal mammalian ovary. *Nature* **428** 145-150.
- 653 **Kelsey TW, Anderson RA, Wright P, Nelson SM & Wallace WH** 2012 Data-driven
654 assessment of the human ovarian reserve. *Mol Hum Reprod* **18** 79-87.
- 655 **Kerr JB, Brogan L, Myers M, Hutt KJ, Mladenovska T, Ricardo S, Hamza K, Scott**
656 **CL, Strasser A & Findlay JK** 2012a The primordial follicle reserve is not
657 renewed after chemical or gamma-irradiation mediated depletion. *Reproduction*
658 **143** 469-476.
- 659 **Kerr JB, Duckett R, Myers M, Britt KL, Mladenovska T & Findlay JK** 2006
660 Quantification of healthy follicles in the neonatal and adult mouse ovary:
661 evidence for maintenance of primordial follicle supply. *Reproduction* **132** 95-109.
- 662 **Kerr JB, Hutt KJ, Michalak EM, Cook M, Vandenberg CJ, Liew SH, Bouillet P,**
663 **Mills A, Scott CL, Findlay JK & Strasser A** 2012b DNA damage-induced
664 primordial follicle oocyte apoptosis and loss of fertility require TAp63-mediated
665 induction of Puma and Noxa. *Mol Cell* **48** 343-352.
- 666 **Kezele P, Nilsson E & Skinner MK** 2002 Cell-cell interactions in primordial follicle
667 assembly and development. *Front Biosci* **7** d1990-1996.
- 668 **Kim JY** 2012 Control of ovarian primordial follicle activation. *Clin Exp Reprod Med* **39**
669 10-14.
- 670 **Kimura F, Bonomi LM & Schneyer AL** 2011 Follistatin regulates germ cell nest
671 breakdown and primordial follicle formation. *Endocrinology* **152** 697-706.
- 672 **Kujjo LL, Chang EA, Pereira RJ, Dhar S, Marrero-Rosado B, Sengupta S, Wang**
673 **H, Cibelli JB & Perez GI** 2011 Chemotherapy-induced late transgenerational
674 effects in mice. *PLoS One* **6** e17877.
- 675 **Kurilo LF** 1981 Oogonia degeneration in human pre- and perinatal oogenesis. *Tsitol*
676 *Genet* **15** 78-82.
- 677 **Lee H-J, Selesniemi K, Niikura Y, Niikura T, Klein R, Dombkowski DM & Tilly JL**
678 **2007** Bone marrow transplantation generates immature oocytes and rescues long-
679 term fertility in a preclinical mouse model of chemotherapy-induced premature
680 ovarian failure. *J Clin Oncol* **25** 3198-3204.
- 681 **Lei L & Spradling AC** 2013 Female mice lack adult germ-line stem cells but sustain
682 oogenesis using stable primordial follicles. *Proc Nat Acad Sci USA* **110** 8585-
683 8590
- 684
- 685 **Livera G, Petre-Lazar B, Guerquin MJ, Trautmann E, Coffigny H & Habert R**
686 **2008** p63 null mutation protects mouse oocytes from radio-induced apoptosis.
687 *Reproduction* **135** 3-12.

- Lobascio AM, Klinger FG, Scaldaferri ML, Farini D & De Felici M** 2007 Analysis of programmed cell death in mouse fetal oocytes. *Reproduction* **134** 241-252.
- Maheshwari A & Fowler PA** 2008 Primordial follicular assembly in humans--revisited. *Zygote* **16** 285-296.
- Mamsen LS, Lutterodt MC, Andersen EW, Byskov AG & Andersen CY** 2011 Germ cell numbers in human embryonic and fetal gonads during the first two trimesters of pregnancy: analysis of six published studies. *Hum Reprod* **26** 2140-2145.
- Marcinkiewicz JL, Balchak SK & Morrison LJ** 2002 The involvement of tumor necrosis factor-alpha (TNF) as an intraovarian regulator of oocyte apoptosis in the neonatal rat. *Front Biosci* **7** d1997-2005.
- Martins da Silva SJ, Bayne RA, Cambray N, Hartley PS, McNeilly AS & Anderson RA** 2004 Expression of activin subunits and receptors in the developing human ovary: activin A promotes germ cell survival and proliferation before primordial follicle formation. *Dev Biol* **266** 334-345.
- McClellan KA, Gosden R & Taketo T** 2003 Continuous loss of oocytes throughout meiotic prophase in the normal mouse ovary. *Dev Biol* **258** 334-348.
- Monget P, Bobe J, Gougeon A, Fabre S, Monniaux D & Dalbies-Tran R** 2012 The ovarian reserve in mammals: a functional and evolutionary perspective. *Mol Cell Endocrinol* **356** 2-12.
- Moniruzzaman M & Miyano T** 2010 Growth of primordial oocytes in neonatal and adult mammals. *J Reprod Dev* **56** 559-566.
- Mork L, Maatouk DM, McMahon JA, Guo JJ, Zhang P, McMahon AP & Capel B** 2012 Temporal differences in granulosa cell specification in the ovary reflect distinct follicle fates in mice. *Biol Reprod* **86** 37.
- Nelson SM, Telfer EE & Anderson RA** 2013 The ageing ovary and uterus: new biological insights. *Hum Reprod Update* **19** 67-83.
- Oatley J & Hunt PA** 2012 Of mice and (wo)men: purified oogonial stem cells from mouse and human ovaries. *Biol Reprod* **86** 196.
- Pacchiarotti J, Maki C, Ramos T, Marh J, Howerton K, Wong J, Pham J, Anorve S, Chow YC & Izadyar F** 2010 Differentiation potential of germ line stem cells derived from the postnatal mouse ovary. *Differentiation* **79** 159-170.
- Pangas SA** 2012 Regulation of the ovarian reserve by members of the transforming growth factor beta family. *Mol Reprod Dev* **79** 666-679.
- Pangas SA, Choi Y, Ballow DJ, Zhao Y, Westphal H, Matzuk MM & Rajkovic A** 2006 Oogenesis requires germ cell-specific transcriptional regulators *Sohlh1* and *Lhx8*. *Proc Natl Acad Sci U S A* **103** 8090-8095.
- Paredes A, Romero C, Dissen GA, DeChiara TM, Reichardt L, Cornea A, Ojeda SR & Xu B** 2004 TrkB receptors are required for follicular growth and oocyte survival in the mammalian ovary. *Dev Biol* **267** 430-449.
- Pepling ME** 2006 From primordial germ cell to primordial follicle: mammalian female germ cell development. *Genesis* **44** 622-632.
- Pepling ME & Spradling AC** 2001 Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles. *Dev Biol* **234** 339-351.
- Pepling ME, Sundman EA, Patterson NL, Gephardt GW, Medico L, Jr. & Wilson KI** 2010 Differences in oocyte development and estradiol sensitivity among mouse strains. *Reproduction* **139** 349-357.

- Perez GI, Robles R, Knudson CM, Flaws JA, Korsmeyer SJ & Tilly JL** 1999 Prolongation of ovarian lifespan into advanced chronological age by Bax-deficiency. *Nat Genet* **21** 200-203.
- Pesce M, Klinger FG & De Felici M** 2002 Derivation in culture of primordial germ cells from cells of the mouse epiblast: phenotypic induction and growth control by Bmp4 signalling. *Mech Dev* **112** 15-24.
- Peters H, Byskov AG & Grinsted J** 1978 Follicular growth in fetal and prepubertal ovaries of humans and other primates. *Clin Endocrinol Metab* **7** 469-485.
- Peters H, Levy E & Crone M** 1962 Deoxyribonucleic acid synthesis in oocytes of mouse embryos. *Nature* **195** 915-916.
- Poljicanin A, Vukusic Pusic T, Vukojevic K, Caric A, Vilovic K, Tomic S, Soljic V & Saraga-Babic M** 2012 The expression patterns of pro-apoptotic and anti-apoptotic factors in human fetal and adult ovary. *Acta Histochem*.
- Rajkovic A, Pangas SA, Ballow D, Suzumori N & Matzuk MM** 2004 NOBOX deficiency disrupts early folliculogenesis and oocyte-specific gene expression. *Science* **305** 1157-1159.
- Reddy P, Adhikari D, Zheng W, Liang S, Hamalainen T, Tohonen V, Ogawa W, Noda T, Volarevic S, Huhtaniemi I & Liu K** 2009 PDK1 signaling in oocytes controls reproductive aging and lifespan by manipulating the survival of primordial follicles. *Hum Mol Genet* **18** 2813-2824.
- Reddy P, Zheng W & Liu K** 2010 Mechanisms maintaining the dormancy and survival of mammalian primordial follicles. *Trends Endocrinol Metab* **21** 96-103.
- Reizel Y, Itzkovitz S, Adar R, Elbaz J, Jinich A, Chapal-Ilani N, Maruvka YE, Nevo N, Marx Z, Horovitz I, Wasserstrom A, Mayo A, Shur I, Benayahu D, Skorecki K, Segal E, Dekel N & Shapiro E** 2012 Cell lineage analysis of the mammalian female germline. *PLoS Genet* **8** e1002477.
- Rodrigues P, Limback D, McGinnis LK, Plancha CE & Albertini DF** 2009 Multiple mechanisms of germ cell loss in the perinatal mouse ovary. *Reproduction* **137** 709-720.
- Rubinstein AD & Kimchi A** 2012 Life in the balance - a mechanistic view of the crosstalk between autophagy and apoptosis. *J Cell Sci* **125** 5259-5268.
- Rucker EB, 3rd, Dierisseau P, Wagner KU, Garrett L, Wynshaw-Boris A, Flaws JA & Hennighausen L** 2000 Bcl-x and Bax regulate mouse primordial germ cell survival and apoptosis during embryogenesis. *Mol Endocrinol* **14** 1038-1052.
- Spears N, Molinek MD, Robinson LL, Fulton N, Cameron H, Shimoda K, Telfer EE, Anderson RA & Price DJ** 2003 The role of neurotrophin receptors in female germ-cell survival in mouse and human. *Development* **130** 5481-5491.
- Spradling AC** 2004 Stem cells: more like a man. *Nature* **428** 133-134.
- Stoop H, Honecker F, Cools M, de Krijger R, Bokemeyer C & Looijenga LH** 2005 Differentiation and development of human female germ cells during prenatal gonadogenesis: an immunohistochemical study. *Hum Reprod* **20** 1466-1476.
- Suh EK, Yang A, Kettenbach A, Bamberger C, Michaelis AH, Zhu Z, Elvin JA, Bronson RT, Crum CP & McKeon F** 2006 p63 protects the female germ line during meiotic arrest. *Nature* **444** 624-628.
- Tingen C, Kim A & Woodruff TK** 2009 The primordial pool of follicles and nest breakdown in mammalian ovaries. *Mol Hum Reprod* **15** 795-803.

- Vandenabeele P, Galluzzi L, Vanden Berghe T & Kroemer G** 2010 Molecular mechanisms of necroptosis: an ordered cellular explosion. *Nat Rev Mol Cell Biol* **11** 700-714.
- Vaskivuo TE, Anttonen M, Herva R, Billig H, Dorland M, te Velde ER, Stenback F, Heikinheimo M & Tapanainen JS** 2001 Survival of human ovarian follicles from fetal to adult life: apoptosis, apoptosis-related proteins, and transcription factor GATA-4. *J Clin Endocrinol Metab* **86** 3421-3429.
- Wallace WH & Kelsey TW** 2010 Human ovarian reserve from conception to the menopause. *PLoS One* **5** e8772.
- Wartenberg H, Ihmer A, Schwarz S, Miething A & Viebahn C** 2001 Mitotic arrest of female germ cells during prenatal oogenesis. A colcemid-like, non-apoptotic cell death. *Anat Embryol (Berl)* **204** 421-435.
- Wen J, Zhang H, Li G, Mao G, Chen X, Wang J, Guo M, Mu X, Ouyang H, Zhang M & Xia G** 2009 PAR6, a potential marker for the germ cells selected to form primordial follicles in mouse ovary. *PLoS One* **4** e7372.
- White YA, Woods DC, Takai Y, Ishihara O, Seki H & Tilly JL** 2012 Oocyte formation by mitotically active germ cells purified from ovaries of reproductive-age women. *Nat Med* **18** 413-421.
- Woods DC, Telfer EE & Tilly JL** 2012 Oocyte family trees: old branches or new stems? *PLoS Genet* **8** e1002848.
- Woods DC, White YA & Tilly JL** 2013 Purification of oogonial stem cells from adult mouse and human ovaries: an assessment of the literature and a view toward the future. *Reprod Sci* **20** 7-15.
- Xu B, Hua J, Zhang Y, Jiang X, Zhang H, Ma T, Zheng W, Sun R, Shen W, Sha J, Cooke HJ & Shi Q** 2011 Proliferating cell nuclear antigen (PCNA) regulates primordial follicle assembly by promoting apoptosis of oocytes in fetal and neonatal mouse ovaries. *PLoS One* **6** e16046.
- Zhang YS, Lu ZY, Yu Y, Li XR, Li WB, Wang YN & Geng Y** 2012 Derivation, culture and retinal pigment epithelial differentiation of human embryonic stem cells using human fibroblast feeder cells. *J Assist Reprod Genet* **29** 735-744.
- Zheng W, Nagaraju G, Liu Z & Liu K** 2012 Functional roles of the phosphatidylinositol 3-kinases (PI3Ks) signaling in the mammalian ovary. *Mol Cell Endocrinol* **356** 24-30.
- Zou K, Yuan Z, Yang Z, Luo H, Sun K, Zhou L, Xiang J, Shi L, Yu Q, Zhang Y, Hou R & Wu J** 2009 Production of offspring from a germline stem cell line derived from neonatal ovaries. *Nat Cell Biol* **11** 631-636.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

